

FREE ARAKI-WOODS FACTORS AND CONNES' BICENTRALIZER PROBLEM

CYRIL HOUDAYER

ABSTRACT. We show that for any type III₁ free Araki-Woods factor $\mathcal{M} = \Gamma(H_{\mathbf{R}}, U_t)''$, the bicentralizer of the free quasi-free state φ_U is trivial. Using Haagerup's Theorem, it follows that there always exists a faithful normal state ψ on $\mathcal{M}$ such that $(\mathcal{M}^\psi)' \cap \mathcal{M} = \mathbf{C}$.

1. INTRODUCTION

Let $\mathcal{M}$ be a separable type III₁ factor and let φ be a faithful normal state on $\mathcal{M}$. For any $x, y \in \mathcal{M}$, set $[x, y] = xy - yx$ and $[x, \varphi] = x\varphi - \varphi x$. The *asymptotic centralizer* of φ is defined by

$$\text{AC}(\varphi) := \{(x_n) \in \ell^\infty(\mathbf{N}, \mathcal{M}) : \|[x_n, \varphi]\| \rightarrow 0\}.$$

Note that $\text{AC}(\varphi)$ is a unital C^* -subalgebra of $\ell^\infty(\mathbf{N}, \mathcal{M})$. The *bicentralizer* of φ is defined by

$$\text{AB}(\varphi) := \{a \in \mathcal{M} : [a, x_n] \rightarrow 0 \text{ ultrastrongly}, \forall (x_n) \in \text{AC}(\varphi)\}.$$

It is well-known that $\text{AB}(\varphi)$ is a von Neumann subalgebra of $\mathcal{M}$, globally invariant under the modular group (σ_t^φ) . Moreover, $\text{AB}(\varphi) \subset (\mathcal{M}^\varphi)' \cap \mathcal{M}$. If $\text{AB}(\varphi) = \mathbf{C}$, it follows from Connes-Størmer Transitivity Theorem ([3]), that $\text{AB}(\psi) = \mathbf{C}$ for any faithful normal state ψ on $\mathcal{M}$. We shall say in this case that $\mathcal{M}$ has *trivial bicentralizer*. Connes conjectured that *any* separable type III₁ factor should have trivial bicentralizer. If there exists a faithful normal state φ on $\mathcal{M}$ such that $(\mathcal{M}^\varphi)' \cap \mathcal{M} = \mathbf{C}$, then $\mathcal{M}$ has trivial bicentralizer. Haagerup proved in [7] that the converse holds true. Haagerup's Theorem lead to the uniqueness of the amenable type III₁ factor (see [2]). The following type III₁ factors are known to have trivial bicentralizer:

- (1) The unique amenable III₁ factor (Haagerup, [7]).
- (2) Full factors that have almost periodic states (Connes, [4]).
- (3) Free products $(\mathcal{M}_1, \varphi_1) * (\mathcal{M}_2, \varphi_2)$ such that the centralizers $\mathcal{M}_i^{\varphi_i}$ have enough unitaries (Barnett, [1]).

In this paper, we show that the bicentralizer is trivial for a large class of type III₁ factors, namely the *free Araki-Woods factors* of Shlyakhtenko

2000 *Mathematics Subject Classification.* 46L10; 46L54.

Key words and phrases. Free Araki-Woods factors; Connes' bicentralizer problem.

([14]). We briefly recall the construction here, see Section 2 for more details. To each real separable Hilbert space $H_{\mathbf{R}}$ together with an orthogonal representation (U_t) of $\mathbf{R}$ on $H_{\mathbf{R}}$, one can associate a von Neumann algebra denoted by $\Gamma(H_{\mathbf{R}}, U_t)''$, called the free Araki-Woods von Neumann algebra. This is the free analog of the factors coming from the CAR relations. The von Neumann algebra $\Gamma(H_{\mathbf{R}}, U_t)''$ comes equipped with a unique *free quasi-free state* denoted by φ_U , which is always normal and faithful on $\Gamma(H_{\mathbf{R}}, U_t)''$. If $\dim H_{\mathbf{R}} \geq 2$, then $\Gamma(H_{\mathbf{R}}, U_t)''$ is a full factor. It is of type III_1 when (U_t) is non-periodic and non-trivial. If the representation (U_t) is almost periodic, then φ_U is an almost periodic state and it follows from [14] that the relative commutant of the centralizer of the free quasi-free state is trivial, i.e. if $\mathcal{M} := \Gamma(H_{\mathbf{R}}, U_t)''$, then $(\mathcal{M}^{\varphi_U})' \cap \mathcal{M} = \mathbf{C}$. In the almost periodic case, results of [6] yield $\mathcal{M}^{\varphi_U} \simeq L(\mathbf{F}_{\infty})$.

When the representation (U_t) has no eigenvectors (e.g. $U_t = \lambda_t$, the left regular representation of $\mathbf{R}$ on $L^2(\mathbf{R}, \mathbf{R})$), then the centralizer $\mathcal{M}^{\varphi_U}$ is trivial. It was unknown in general whether or not $\Gamma(H_{\mathbf{R}}, U_t)''$ had trivial bicentralizer. Even though the centralizer of the free quasi-free state φ_U may be trivial, we will show that the bicentralizer of φ_U is always trivial. The main result of this paper is the following:

Theorem. *Let $\mathcal{M} := \Gamma(H_{\mathbf{R}}, U_t)''$ be a free Araki-Woods factor of type III_1 . Denote by φ_U the free quasi-free state. Then $\text{AB}(\varphi_U) = \mathbf{C}$. Consequently, there always exists a faithful normal state ψ on $\mathcal{M}$ such that $(\mathcal{M}^{\psi})' \cap \mathcal{M} = \mathbf{C}$.*

Acknowledgements. Part of this work was done while the author was visiting the University of Tokyo. He gratefully thanks Professors Y. Kawahigashi and N. Ozawa for their kind invitation.

2. PRELIMINARIES

2.1. Preliminaries on Spectral Analysis. We shall need a few definitions and results from the spectral theory of abelian automorphism groups. Let (α_t) be an ultraweakly continuous one-parameter automorphism group on a von Neumann algebra $\mathcal{M}$. For $f \in L^1(\mathbf{R})$ and $x \in \mathcal{M}$, set

$$\alpha_f(x) = \int_{-\infty}^{+\infty} f(t) \alpha_t(x) dt.$$

The α -spectrum $\text{Sp}_{\alpha}(x)$ of $x \in \mathcal{M}$ is defined as the set of characters $\gamma \in \widehat{\mathbf{R}}$ for which $\widehat{f}(\gamma) = 0$, for all $f \in L^1(\mathbf{R})$ satisfying $\alpha_f(x) = 0$. We shall identify $\widehat{\mathbf{R}}$ with $\mathbf{R}$ in the usual way, such that

$$\widehat{f}(\gamma) = \int_{-\infty}^{+\infty} e^{i\gamma t} f(t) dt, \forall \gamma \in \mathbf{R}, \forall f \in L^1(\mathbf{R}).$$

For $z \in \mathbf{C}$, denote by $\Im(z)$ its imaginary part.

Lemma 2.1 ([7]). *Let $\mathcal{M}$ and (α_t) be as above. Let $x \in \mathcal{M}$ and $\delta > 0$. If the function $t \mapsto \alpha_t(x)$ can be extended to an entire (analytic) $\mathcal{M}$ -valued function such that*

$$\|\alpha_z(x)\| \leq Ce^{\delta|\Im(z)|}, \forall z \in \mathbf{C},$$

for some constant $C > 0$, then $\text{Sp}_\alpha(x) \subset [-\delta, \delta]$.

Let φ be a f.n. state on a von Neumann algebra $\mathcal{M}$. Denote by (σ_t^φ) the modular group on $\mathcal{M}$ of the state φ . Denote by $L^2(\mathcal{M}, \varphi)$ the L^2 -space associated with φ and by ξ_φ the canonical cyclic separating vector. We shall write $\|x\|_\varphi = \varphi(x^*x)^{1/2}$, for any $x \in \mathcal{M}$. On bounded subsets of $\mathcal{M}$, the topology given by the norm $\|\cdot\|_\varphi$ coincides with the strong-operator topology. Recall that $S_\varphi^0 : x\xi_\varphi \mapsto x^*\xi_\varphi$ is a closable (densely defined) operator on $L^2(\mathcal{M}, \varphi)$. Denote by S_φ its closure and write $S_\varphi = J_\varphi \Delta_\varphi^{1/2}$ for its polar decomposition. Note that $L^2(\mathcal{M}, \varphi)$ is naturally endowed with an $\mathcal{M}$ - $\mathcal{M}$ bimodule structure defined as follows:

$$\begin{aligned} x \cdot \xi &:= x\xi \\ \xi \cdot x &:= J_\varphi x^* J_\varphi \xi, \forall x \in \mathcal{M}, \forall \xi \in L^2(\mathcal{M}, \varphi). \end{aligned}$$

We shall simply denote $x \cdot \xi$ and $\xi \cdot x$ by $x\xi$ and ξx . The next lemma is well-known, but we give a proof for the reader's convenience.

Lemma 2.2. *Let $\mathcal{M}$ and φ be as above. Let $x \in \mathcal{M}$ and $0 < \delta < 1$. Assume that $\text{Sp}_{\sigma^\varphi}(x) \subset [-\delta, \delta]$. Then $\|x\xi_\varphi - \xi_\varphi x\| \leq \delta\|x\|_\varphi$.*

Proof. Let $x \in \mathcal{M}$ and $0 < \delta < 1$ such that $\text{Sp}_{\sigma^\varphi}(x) \subset [-\delta, \delta]$. Let $f \in L^1(\mathbf{R})$ for which the Fourier transform $\hat{f}$ vanishes on $[-\delta, \delta]$. Since $\text{Sp}_{\sigma^\varphi}(\sigma_f^\varphi(x)) \subset \text{Sp}_{\sigma^\varphi}(x) \cap \text{support}(\hat{f}) = \emptyset$ (see [5]), it follows that $\sigma_f^\varphi(x) = 0$. We have

$$\begin{aligned} \hat{f}(\log \Delta_\varphi) x \xi_\varphi &= \int_{-\infty}^{+\infty} f(t) \Delta_\varphi^{it} x \xi_\varphi dt \\ &= \int_{-\infty}^{+\infty} f(t) \sigma_t^\varphi(x) \xi_\varphi dt \\ &= \sigma_f^\varphi(x) \xi_\varphi \\ &= 0. \end{aligned}$$

Thus, by approximating $\mathbf{1}_{\mathbf{R} \setminus [-\delta, \delta]}$ by such functions $\hat{f}$, we get

$$\mathbf{1}_{\mathbf{R} \setminus [-\delta, \delta]}(\log \Delta_\varphi) x \xi_\varphi = 0,$$

i.e. $x\xi_\varphi$ is in the spectral subspace of $\log \Delta_\varphi$ corresponding to the interval $[-\delta, \delta]$. Notice that

$$\xi_\varphi x = J_\varphi x^* J_\varphi \xi_\varphi = J_\varphi x^* \xi_\varphi = J_\varphi S_\varphi x \xi_\varphi = \Delta_\varphi^{1/2} x \xi_\varphi.$$

Clearly, $\sup\{|e^{t/2} - 1| : t \in [-\delta, \delta]\} = e^{\delta/2} - 1$. Moreover, the operator $(1 - \Delta_\varphi^{1/2})\mathbf{1}_{[-\delta, \delta]}(\log \Delta_\varphi)$ is bounded and precisely

$$\|(1 - \Delta_\varphi^{1/2})\mathbf{1}_{[-\delta, \delta]}(\log \Delta_\varphi)\| \leq e^{\delta/2} - 1 \leq \delta,$$

since $0 < \delta < 1$. Thus, we get

$$\begin{aligned}
\|x\xi_\varphi - \xi_\varphi x\| &= \|(1 - \Delta_\varphi^{1/2})x\xi_\varphi\| \\
&= \|(1 - \Delta_\varphi^{1/2})\mathbf{1}_{[-\delta, \delta]}(\log \Delta_\varphi)x\xi_\varphi\| \\
&\leq \|(1 - \Delta_\varphi^{1/2})\mathbf{1}_{[-\delta, \delta]}(\log \Delta_\varphi)\| \|x\xi_\varphi\| \\
&\leq \delta \|x\|_\varphi.
\end{aligned}$$

□

Lemma 2.3 ([7]). *Let $\mathcal{M}$ and φ be as above. Let $(x_n) \in \ell^\infty(\mathbf{N}, \mathcal{M})$. Then*

$$\lim_n \|x_n \xi_\varphi - \xi_\varphi x_n\| = 0 \iff \lim_n \|x_n \varphi - \varphi x_n\| = 0.$$

2.2. Preliminaries on Shlyakhtenko's Free Araki-Woods Factors.

Recall now the construction of the free Araki-Woods factors due to Shlyakhtenko ([14]). Let $H_{\mathbf{R}}$ be a real separable Hilbert space and let (U_t) be an orthogonal representation of $\mathbf{R}$ on $H_{\mathbf{R}}$. Let $H = H_{\mathbf{R}} \otimes_{\mathbf{R}} \mathbf{C}$ be the complexified Hilbert space. Let J be the canonical anti-unitary involution on H defined by:

$$J(\xi + i\eta) = \xi - i\eta, \forall \xi, \eta \in H_{\mathbf{R}}.$$

If A is the infinitesimal generator of (U_t) on H , we recall that $j : H_{\mathbf{R}} \rightarrow H$ defined by $j(\zeta) = (\frac{2}{A^{-1}+1})^{1/2}\zeta$ is an isometric embedding of $H_{\mathbf{R}}$ into H . Moreover $JAJ = A^{-1}$ and $JA^{it} = A^{it}J$, for every $t \in \mathbf{R}$. Let $K_{\mathbf{R}} = j(H_{\mathbf{R}})$. It is easy to see that $K_{\mathbf{R}} \cap iK_{\mathbf{R}} = \{0\}$ and $K_{\mathbf{R}} + iK_{\mathbf{R}}$ is dense in H . Write $T = JA^{-1/2}$. Then T is an anti-linear closed invertible operator on H satisfying $T = T^{-1}$. Such an operator is called an *involution* on H . Moreover, $K_{\mathbf{R}} = \{\xi \in \text{dom}(T) : T\xi = \xi\}$.

We introduce the *full Fock space* of H :

$$\mathcal{F}(H) = \mathbf{C}\Omega \oplus \bigoplus_{n=1}^{\infty} H^{\otimes n}.$$

The unit vector Ω is called the *vacuum vector*. For any $\xi \in H$, define the *left creation operator*

$$\ell(\xi) : \mathcal{F}(H) \rightarrow \mathcal{F}(H) : \begin{cases} \ell(\xi)\Omega = \xi, \\ \ell(\xi)(\xi_1 \otimes \cdots \otimes \xi_n) = \xi \otimes \xi_1 \otimes \cdots \otimes \xi_n. \end{cases}$$

We have $\|\ell(\xi)\| = \|\xi\|$ and $\ell(\xi)$ is an isometry if $\|\xi\| = 1$. For any $\xi \in H$, we denote by $s(\xi)$ the real part of $\ell(\xi)$ given by

$$s(\xi) = \frac{\ell(\xi) + \ell(\xi)^*}{2}.$$

The crucial result of Voiculescu [16] claims that the distribution of the operator $s(\xi)$ with respect to the vacuum vector state $\varphi(x) = \langle x\Omega, \Omega \rangle$ is the semicircular law of Wigner supported on the interval $[-\|\xi\|, \|\xi\|]$.

Definition 2.4 (Shlyakhtenko, [14]). Let (U_t) be an orthogonal representation of $\mathbf{R}$ on the real Hilbert space $H_{\mathbf{R}}$. The *free Araki-Woods* von Neumann algebra associated with $(H_{\mathbf{R}}, U_t)$, denoted by $\Gamma(H_{\mathbf{R}}, U_t)''$, is defined by

$$\Gamma(H_{\mathbf{R}}, U_t)'' := \{s(\xi) : \xi \in K_{\mathbf{R}}\}''.$$

The vector state $\varphi_U(x) = \langle x\Omega, \Omega \rangle$ is called the *free quasi-free state* and is faithful on $\Gamma(H_{\mathbf{R}}, U_t)''$. Let $\xi, \eta \in K_{\mathbf{R}}$ and write $\zeta = \xi + i\eta$. We have

$$2s(\xi) + 2is(\eta) = \ell(\zeta) + \ell(T\zeta)^*.$$

Thus, $\Gamma(H_{\mathbf{R}}, U_t)''$ is generated as a von Neumann algebra by the operators of the form $\ell(\zeta) + \ell(T\zeta)^*$ where $\zeta \in \text{dom}(T)$. Note that the modular group $(\sigma_t^{\varphi_U})$ of the free quasi-free state φ_U is given by $\sigma_{-t}^{\varphi_U} = \text{Ad}(\mathcal{F}(U_t))$, where $\mathcal{F}(U_t) = \text{id} \oplus \bigoplus_{n \geq 1} U_t^{\otimes n}$. In particular, it satisfies

$$\sigma_{-t}^{\varphi_U}(\ell(\zeta) + \ell(T\zeta)^*) = \ell(U_t\zeta) + \ell(TU_t\zeta)^*, \forall \zeta \in \text{dom}(T), \forall t \in \mathbf{R}.$$

The free Araki-Woods factors provided many new examples of full factors of type III [1, 5, 11]. We can summarize the general properties of the free Araki-Woods factors in the following theorem (see also [15]):

Theorem 2.5 (Shlyakhtenko, [11, 12, 13, 14]). *Let (U_t) be an orthogonal representation of $\mathbf{R}$ on the real Hilbert space $H_{\mathbf{R}}$ with $\dim H_{\mathbf{R}} \geq 2$. Denote by $\mathcal{M} := \Gamma(H_{\mathbf{R}}, U_t)''$.*

- (1) *$\mathcal{M}$ is a full factor and Connes' invariant $\tau(\mathcal{M})$ is the weakest topology on $\mathbf{R}$ that makes the map $t \mapsto U_t$ strongly continuous.*
- (2) *$\mathcal{M}$ is of type II_1 iff $U_t = \text{id}$, for every $t \in \mathbf{R}$.*
- (3) *$\mathcal{M}$ is of type III_λ ($0 < \lambda < 1$) iff (U_t) is periodic of period $\frac{2\pi}{|\log \lambda|}$.*
- (4) *$\mathcal{M}$ is of type III_1 in the other cases.*
- (5) *The factor $\mathcal{M}$ has almost periodic states iff (U_t) is almost periodic.*

Let $H_{\mathbf{R}} = \mathbf{R}^2$ and $0 < \lambda < 1$. Let

$$(1) \quad U_t^\lambda = \begin{pmatrix} \cos(t \log \lambda) & -\sin(t \log \lambda) \\ \sin(t \log \lambda) & \cos(t \log \lambda) \end{pmatrix}.$$

Notation 2.6 ([14]). Denote by $(T_\lambda, \varphi_\lambda) := (\Gamma(H_{\mathbf{R}}, U_t)'', \varphi_U)$ where $H_{\mathbf{R}} = \mathbf{R}^2$ and (U_t) is given by Equation (1).

Using a powerful tool called the *matricial model*, Shlyakhtenko was able to prove the following isomorphism

$$(T_\lambda, \varphi_\lambda) \cong (\mathbf{B}(\ell^2(\mathbf{N})), \psi_\lambda) * (L^\infty[-1, 1], \mu),$$

where $\psi_\lambda(e_{ij}) = \delta_{ij}\lambda^j(1 - \lambda)$, $i, j \in \mathbf{N}$, and μ is a non-atomic measure on $[-1, 1]$. The notation $\cong$ means a state-preserving isomorphism. He also proved that $(T_\lambda, \varphi_\lambda)$ has the *free absorption property*, namely,

$$(T_\lambda, \varphi_\lambda) * L(\mathbf{F}_\infty) \cong (T_\lambda, \varphi_\lambda).$$

3. THE MAIN RESULT

3.1. Technical Lemmas. As we said before, the centralizer of the free quasi-free state may be trivial; this is the case for instance when the orthogonal representation (U_t) on $H_{\mathbf{R}}$ has no eigenvectors. Nevertheless, the following lemma shows that for any free Araki-Woods von Neumann algebra, there exists a non-trivial sequence of unitaries (u_n) in the asymptotic centralizer of the free quasi-free state φ_U .

Lemma 3.1 (Vaes, [15]). *Let $\mathcal{M} := \Gamma(H_{\mathbf{R}}, U_t)''$ be a free Araki-Woods von Neumann algebra. Denote by φ the free quasi-free state and by (σ_t) the modular group of the state φ . Then there exists a sequence of unitaries (u_n) in $\mathcal{M}$ entire (analytic) w.r.t. (σ_t) such that*

- (1) $\|\sigma_z(u_n) - u_n\| \rightarrow 0$ uniformly on compact sets of $\mathbf{C}$;
- (2) $\varphi(u_n) \rightarrow 0$;
- (3) $(u_n) \in \text{AC}(\varphi)$.

Proof. This lemma, with the exception of item (3), is Vaes' result (see Lemma 4.3 in [15]). Item (3) was not observed by Vaes but is immediate from the construction using Lemmas 2.1, 2.2, 2.3. $\square$

The following lemma is a generalization of Barnett's lemma (see [1]) which was itself a generalization of Murray & von Neumann's 14ϵ lemma.

Lemma 3.2 (Vaes, [15]). *For $i = 1, 2$, let $(\mathcal{M}_i, \varphi_i)$ be a von Neumann algebra endowed with a f.n. state. Denote by $(\mathcal{M}, \varphi) = (\mathcal{M}_1, \varphi_1) * (\mathcal{M}_2, \varphi_2)$ the free product. Let $a \in \mathcal{M}_1$, $b, c \in \mathcal{M}_2$. Assume that a, b, c belong to the domain of $\sigma_{i/2}^\varphi$. Then, for every $x \in \mathcal{M}$,*

$$\|x - \varphi(x)1\|_\varphi \leq \mathcal{E}(a, b, c) \max \{ \| [x, a] \|_\varphi, \| [x, b] \|_\varphi, \| [x, c] \|_\varphi \} + \mathcal{F}(a, b, c) \|x\|_\varphi$$

where

$$\begin{aligned} \mathcal{E}(a, b, c) &= 6\|a\|^3 + 4\|b\|^3 + 4\|c\|^3 \\ \mathcal{F}(a, b, c) &= 3\mathcal{C}(a) + 2\mathcal{C}(b) + 2\mathcal{C}(c) + 12|\varphi(cb^*)| \|cb^*\| \\ \mathcal{C}(a) &= 2\|a\|^3 \|\sigma_{i/2}^\varphi(a) - a\| + 2\|a\|^2 \|a^*a - 1\| \\ &\quad + 3(1 + \|a\|^2) \|aa^* - 1\| + 6|\varphi(a)| \|a\|. \end{aligned}$$

3.2. Proof of the Theorem. Let $\mathcal{M} := \Gamma(H_{\mathbf{R}}, U_t)''$ be a free Araki-Woods factor of type III₁ and denote by φ the free quasi-free state. We recall that such a factor can always be written as the free product of three free Araki-Woods von Neumann algebras (see the proof of Theorem 2.7 in [15]):

$$(\mathcal{M}, \varphi) \cong (\mathcal{M}_1, \varphi_1) * (\mathcal{M}_2, \varphi_2) * (\mathcal{M}_3, \varphi_3).$$

Notice that $\sigma_t^\varphi = \sigma_t^{\varphi_1} * \sigma_t^{\varphi_2} * \sigma_t^{\varphi_3}, \forall t \in \mathbf{R}$.

Thanks to Lemma 3.1, we may choose three sequences of unitaries (u_n^j) , for $j \in \{1, 2, 3\}$, such that $u_n^j \in \mathcal{U}(\mathcal{M}_j)$ is analytic w.r.t. $(\sigma_t^{\varphi_j})$, and satisfies conditions (1) – (3) of Lemma 3.1, for all $j \in \{1, 2, 3\}$. The way the sequence

of unitaries (u_n^j) is constructed in Lemma 3.1 (see Lemma 4.3 in [15]) shows that conditions (1) – (3) are satisfied for the state φ , i.e. the sequence of unitaries (u_n^j) in $\mathcal{M}_j$ satisfies for every $j \in \{1, 2, 3\}$,

- (1) $\|\sigma_z^\varphi(u_n^j) - u_n^j\| \rightarrow 0$, uniformly on compact sets of $\mathbf{C}$;
- (2) $\varphi(u_n^j) \rightarrow 0$;
- (3) $\|[u_n^j, \varphi]\| \rightarrow 0$.

Moreover, by freeness, $\varphi(u_n^3(u_n^2)^*) = \varphi(u_n^3)\overline{\varphi(u_n^2)} \rightarrow 0$.

Assume $a \in \text{AB}(\varphi)$. Fix $\varepsilon > 0$. Since $(u_n^j) \in \text{AC}(\varphi)$, it follows that $[a, u_n^j] \rightarrow 0$ ultrastrongly for any $j \in \{1, 2, 3\}$, and thus we may choose $n \in \mathbf{N}$ large enough such that

$$\begin{aligned} \|[a, u_n^j]\|_\varphi &\leq \varepsilon/28, \forall j \in \{1, 2, 3\}, \\ \mathcal{F}(u_n^1, u_n^2, u_n^3)\|a\|_\varphi &\leq \varepsilon/2. \end{aligned}$$

Thus, we get thanks to Lemma 3.2, $\|a - \varphi(a)1\|_\varphi \leq \varepsilon/2 + \varepsilon/2 = \varepsilon$. Since $\varepsilon > 0$ is arbitrary, $a = \varphi(a)1$. Thus $\text{AB}(\varphi) = \mathbf{C}$ and we are done.

3.3. Final Remark. Set $\mathcal{M} := \Gamma(L^2(\mathbf{R}, \mathbf{R}), \lambda_t)''$ the free Araki-Woods factor associated with the left regular representation (λ_t) of $\mathbf{R}$ on the real Hilbert space $L^2(\mathbf{R}, \mathbf{R})$. Shlyakhtenko showed in [13] that the continuous core of $\mathcal{M}$ is isomorphic to $L(\mathbf{F}_\infty) \otimes \mathbf{B}(\ell^2)$ and the dual action is precisely the one constructed by Rădulescu in [9]. As noticed in [10], for any f.n. state φ on $\mathcal{M}$, the centralizer $\mathcal{M}^\varphi$ is amenable. Indeed, first we have

$$(2) \quad \mathcal{M} \rtimes_{\sigma^\varphi} \mathbf{R} \simeq L(\mathbf{F}_\infty) \otimes \mathbf{B}(\ell^2).$$

Choose on the left hand-side of (2) a non-zero projection $p \in L(\mathbf{R})$ such that $\text{Tr}(p) < +\infty$. We know that $p(\mathcal{M} \rtimes_{\sigma^\varphi} \mathbf{R})p \simeq L(\mathbf{F}_\infty)$ is *solid* by Ozawa's result ([8]). Since $L(\mathbf{R})p$ is diffuse in $p(\mathcal{M} \rtimes_{\sigma^\varphi} \mathbf{R})p$, its relative commutant must be amenable. In particular $\mathcal{M}^\varphi \otimes L(\mathbf{R})p$ is amenable. Thus, $\mathcal{M}^\varphi$ is amenable. Consequently, we obtain

Corollary 3.3. *Let $\mathcal{M} := \Gamma(L^2(\mathbf{R}, \mathbf{R}), \lambda_t)''$. Then there exists a f.n. state ψ on $\mathcal{M}$ such that $(\mathcal{M}^\psi)' \cap \mathcal{M} = \mathbf{C}$. Moreover, $\mathcal{M}^\psi$ is isomorphic to the unique hyperfinite II_1 factor.*

REFERENCES

- [1] L. BARNETT, *Free product von Neumann algebras of type III*. Proc. Amer. Math. Soc. **123** (1995), 543–553.
- [2] A. CONNES, *Factors of type III_1 , property L'_λ and closure of inner automorphisms*. J. Operator Theory **14** (1985), 189–211.
- [3] A. CONNES & E. STÖRMER, *Homogeneity of the state space of factors of type III_1* . J. Funct. Anal. **28** (1978), 187–196.
- [4] A. CONNES, *Almost periodic states and factors of type III_1* . J. Funct. Anal. **16** (1974), 415–445.
- [5] A. CONNES, *Une classification des facteurs de type III*. Ann. Sci. École Norm. Sup. **6** (1973), 133–252.

- [6] K. DYKEMA, *Free products of finite-dimensional and other von Neumann algebras with respect to non-tracial states*. Free probability theory (Waterloo, ON, 1995) Fields Inst. Commun. **12** Amer. Math. Soc., Providence, RI, 1997, pp. 41–88.
- [7] U. HAAGERUP, *Connes' bicentralizer problem and uniqueness of the injective factor of type III₁*. Acta Math. **69** (1986), 95–148.
- [8] N. OZAWA, *Solid von Neumann algebras*. Acta Math. **192** (2004), 111–117.
- [9] F. RĂDULESCU, *A one-parameter group of automorphisms of $L(\mathbf{F}_\infty) \otimes \mathbf{B}(H)$ scaling the trace*. C. R. Acad. Sci. Paris Sr. I Math. **314** (1992), 1027–1032.
- [10] D. SHLYAKHTENKO, *On multiplicity and free absorption for free Araki-Woods factors*. math.OA/0302217
- [11] D. SHLYAKHTENKO, *On the classification of full factors of type III*. Trans. Amer. Math. Soc. **356** (2004), 4143–4159.
- [12] D. SHLYAKHTENKO, *A-valued semicircular systems*. J. Funct. Anal. **166** (1999), 1–47.
- [13] D. SHLYAKHTENKO, *Some applications of freeness with amalgamation*. J. reine angew. Math. **500** (1998), 191–212.
- [14] D. SHLYAKHTENKO, *Free quasi-free states*. Pacific J. Math. **177** (1997), 329–368.
- [15] S. VAES, *États quasi-libres libres et facteurs de type III (d'après D. Shlyakhtenko)*. Séminaire Bourbaki, exposé 937, Astérisque **299** (2005), 329–350.
- [16] D.-V. VOICULESCU, K.J. DYKEMA & A. NICA, *Free random variables*. CRM Monograph Series **1**. American Mathematical Society, Providence, RI, 1992.

CNRS-ENS LYON, UMPA UMR 5669, 69364 LYON CEDEX 7, FRANCE

E-mail address: `cyril.houdayer@umpa.ens-lyon.fr`